

**UNCLASSIFIED**

---

**AD 273 978**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

ASTIA


273978

AD NO.

AD NO.

MASS SPECTROMETRIC STUDY OF THE  
VAPORIZATION OF THE  
Ti-B SYSTEM

UNION  
CARBIDE



PARMA RESEARCH LABORATORY  
UNION CARBIDE CORPORATION

77

Research Report  
No. C-11

**Mass Spectrometric Study of the Vaporization  
of the Ti-B System**

**P. O. Schissel  
Parma Research Laboratory  
Union Carbide Corporation  
Parma, Ohio**

**O. C. Trulson  
Union Carbide Research Institute  
Tarrytown, New York**

**February 5, 1962**

**Under the Auspices of the  
Advanced Research Projects Agency  
and  
Army Rocket and Guided Missile Agency  
through Contract No. DA-30-069-ORD-2787  
with the Union Carbide Research Institute  
Box 278, Tarrytown, New York**

MASS SPECTROMETRIC STUDY OF THE VAPORIZATION  
OF THE Ti-B SYSTEM\*

P. O. Schissel

and

O. C. Trulson\*\*

I. ABSTRACT

A mass spectrometer has been used with Knudsen cells to study the vaporization of the titanium-boron system. The pressures of Ti(g) and B(g) have been determined over several condensed phases and yield  $\Delta H_{298}^{\circ} = 430$  kcal/mole and  $\Delta H_{298}^{\circ} = -52$  kcal/mole for the heats of vaporization and formation of TiB<sub>2</sub>, respectively. Thermodynamic functions for TiB(s) have been obtained.

II. INTRODUCTION

Brewer and Haraldsen<sup>1</sup> have summarized thermodynamic information on some refractory borides and have made qualitative estimates of heats of formation based on the relative stabilities of boron-containing compounds at high temperatures. They mention the experimental difficulties associated with borides and suggest vapor pressure measurements as a promising approach to the evaluation of thermodynamic quantities. Vapor pressure measurements on elemental boron

---

\*This work was supported by the Advanced Research Projects Agency.

\*\*Union Carbide Research Institute.

have been made by Chupka<sup>2</sup>, Thorn<sup>2</sup>, Schissel and Williams<sup>3</sup>, and Akishin et al<sup>4</sup>, who find values in agreement for the heat of sublimation near  $\Delta H_0^0 = 130$  kcal/mole.

The qualitative observations of Brewer and Haraldsen<sup>1</sup> indicate a heat of formation for  $TiB_2(s)$  near -72 kcal/mole; this value is confirmed by Samsonov.<sup>5</sup> However, mass spectrometric determinations of partial vapor pressures by Schissel and Williams<sup>3</sup> showed  $TiB_2$  to be more volatile than would correspond to  $\Delta H_f^0 = -72$  kcal/mole, leading to a value of about -32 kcal/mole instead; the congruent vaporization of the Ti-B system as used by Searcy et al<sup>6</sup> provided bounds for the relative electron impact ionization cross sections for titanium and boron vapor and substantiated the conclusions of Schissel and Williams; and a high temperature calorimetric measurement by Lowell and Williams<sup>7</sup> resulted in  $\Delta H_f = -50 \pm 5$  kcal/mole.

A mass spectrometer was employed in the present work with Knudsen effusion cells where the pressures of titanium and boron vapor were determined from rates of effusion. The vaporization of  $TiB_2$  was studied under several experimental conditions: (1)  $TiB_2$  with excess boron added, (2)  $TiB_2$ , and (3)  $TiB_2$  with excess titanium added. The results of each set of measurements were used to compute the heat of vaporization of  $TiB_2$ , and current thermodynamic data were used to obtain the heat of formation at 298°K.

### III. APPARATUS

The mass spectrometer, similar to that developed by Inghram and co-workers<sup>8</sup>, was a 60° sector instrument of 12 inch radius, differentially pumped

by two 5-liter/sec ion-getter vacuum pumps.\* Electron bombardment was used to heat the Knudsen cells in an arrangement similar to that of Chupka and Inghram.<sup>9</sup> A movable shutter interposed between the cell and the electron-impact ionization source was used to distinguish neutral species from background non-condensable molecules of the same mass. The energy of the ionizing electrons was continuously adjustable from 5 to 150 ev for appearance potential measurements. The main detector\*\* was a twenty-stage electron multiplier employing Be-Cu dynodes. Provision was also made to collect the ion beam directly on an adjustable plate which could be moved in and out of the line of the ion beam. The output from each collector was measured with an electrometer and recorded continuously on a strip-chart recorder.

Temperature measurements were made with a Leeds and Northrup disappearing-filament optical pyrometer sighted into small holes (0.030"dia, 0.060" deep) in the sides of the Knudsen cells. The pyrometer was calibrated against a standard lamp certified by the National Bureau of Standards to be accurate to  $\pm 7^{\circ}\text{C}$  at the highest temperatures employed in this study. An auxiliary check was made against a pyrometer calibrated by the National Bureau of Standards and no systematic differences were noted. The calibration error was therefore taken to be  $\pm 7^{\circ}\text{C}$ . The temperature correction for the sight window was determined before and after all data were taken. The estimated total error was  $\pm 20^{\circ}$  for the work with boron added to  $\text{TiB}_2$  and  $\pm 10^{\circ}\text{C}$  for the remaining measurements. Because of the high emissivity of graphite and since the temperatures were measured by sighting in holes in the graphite, no emissivity corrections were necessary. Further tests on

---

\* Varian Associates, 611 Hansen Way, Palo Alto, California.

\*\* Nuclide Analysis Associates, Box 752, State College, Pennsylvania.

the temperature measurements showed the above statement to be correct and further that temperature gradients in the cell were small. In an auxiliary vacuum system two pyrometers were simultaneously sighted on a tungsten ribbon filament to obtain their relative calibration. The Knudsen cell was then placed at the same position as the tungsten ribbon with one pyrometer sighted into the temperature measuring hole in the side of the cell and the second pyrometer sighted into the Knudsen effusion hole. At the temperatures used in the experiments for which Third Law calculations were made in this study no systematic difference could be measured, but at the very lowest temperatures, (1700°K), attained only during measurements pertaining to the Second Law treatment of the vaporization of boron, a temperature difference was noted but never exceeded 14°C.

#### IV. PROCEDURE AND RESULTS

The phase diagram shown by Brewer and Haraldsen<sup>1</sup> for the Ti-B-C system indicates the expected mutual stabilities of the various phases. This diagram was used to determine suitable crucible materials for the systems which were studied. For the experimental situations discussed below the phase systems are identified by the sample materials initially loaded in the crucible. Subsequent X-ray analyses of the samples after heating showed partial conversion to other phases, particularly in the boron and titanium rich cases. The experimental results are discussed in the light of the X-ray observations made at room temperatures after the runs. In this paper all solid-solid reactions are considered in the approximation  $\Delta C_p = 0$  and therefore phases stable over limited temperature ranges are ignored.



TiB<sub>2</sub>(s) + B(s) — X-ray analysis of an aliquot part of the TiB<sub>2</sub> and boron powdered samples showed no detectable impurities, and spectroscopic analysis showed 0.1% iron, 0.1% silicon and smaller amounts of chromium and other metals. The samples were run in a B<sub>4</sub>C crucible (11/32" diameter, 11/16" long, 1/32" wall thickness) which was contained in a graphite outer jacket (15/16" diameter, 1-1/4" long, 1/8" wall thickness). Two B<sub>4</sub>C crucibles were made by heating graphite crucibles in the presence of boron. The first crucible was used to demonstrate the complete boronization of the graphite; the second was used for the experimental runs. Both crucible and jacket had rectangular orifices 4.5 mm long and 0.75 mm wide which provided a ratio of orifice area to sample area of  $\sigma_A \approx 0.04$  (sample area means gross exposed powder area, not the microscopic area which may make the effective  $\sigma_A$  smaller).

After the sample was loaded, the Knudsen cell was heated slowly to the temperature at which a shutter effect was observed on Ti<sup>48</sup> and B<sup>11</sup>. (These isotopic signals were used for all the measurements, whereas signals at other isotopic positions were used only to confirm the identification of the species). During the preliminary outgassing period, gaseous titanium oxide and boron oxide species were observed but disappeared completely after several hours of heating. No data were taken until signals obviously characteristic of the oxide species had disappeared and appearance-potential measurements indicated that the B and Ti signals resulted solely from the elemental gases.

Data were taken on the temperature dependence of the boron signal and were plotted according to the Clausius-Clapeyron relation to obtain the heat of

sublimation of boron,  $\Delta H_T^0$ . Four experimental runs of this type were made. The results for  $\Delta H_T^0$  in kcal/mole obtained chronologically are:  $130.2 \pm 4.4$  at  $1962^\circ\text{K}$ ,  $128.4 \pm 1.2$  at  $2085^\circ\text{K}$ ,  $133.0 \pm 2.2$  at  $2072^\circ\text{K}$ , and  $132.6 \pm 1.5$  at  $2121^\circ\text{K}$ , where the errors are the statistical errors determined from least square analyses, and the temperatures are the average midrange temperatures of the individual runs. Figure 1 shows a plot of the data of the second run where the relative boron vapor pressure ( $I_B T$ ) is plotted as ordinate against  $1/T$  as abscissa. Tabulated data<sup>2</sup> were used to reduce the  $\Delta H_T^0$  values to  $\Delta H_0^0$  with the results  $\Delta H_0^0 = 130.8 \pm 5.1$ ,  $129.3 \pm 2.7$ ,  $133.8 \pm 3.4$ , and  $133.5 \pm 2.9$  kcal/mole. The final error assignments were obtained from the statistical combination of random errors from the least square analyses and from the instrumental uncertainty in the absolute temperature (fractional error in  $\Delta H_T^0 = 2 \frac{\Delta T}{T}$ ) as described by Trulson, et al<sup>10</sup>. A powder pattern X-ray analysis of the sample after the runs showed strong  $\text{TiB}_2$  lines and no boron lines. While the boron lines were weak even at the start, it is possible that all free boron was lost during the course of heating. Nevertheless, the average of the above Second Law values,  $131.9$  kcal/mole, is only slightly higher than mass spectrometric determinations using the Third Law<sup>2, 3, 4</sup>, and is in excellent agreement with recently tabulated data<sup>2</sup>. Therefore, computations in this section will utilize the tabulated data for the vapor pressure of boron.

The titanium pressure was determined relative to the boron pressure from

$$P_{\text{Ti}} = P_{\text{B}} \frac{I_{\text{Ti}} T_{\text{Ti}}}{I_{\text{B}} T_{\text{B}}} \frac{(\sigma_i S)_{\text{B}}}{(\sigma_i S)_{\text{Ti}}} \quad (1)$$

where P denotes the partial pressure, I the relative collected ion current,  $\sigma$  the cross section for ionization of neutral species, T the temperature, i the isotopic abundance correction factor, and S the electron multiplier conversion factor. The ionization cross sections computed by Otvos and Stevenson<sup>11</sup> were used, and the multiplier conversion efficiencies were determined directly for both boron and titanium signals. The relative ion signals were measured at the same temperature. Electron bombardment energies ranging from 15 to 60 ev were used, but were always normalized to 40 ev by means of appearance-potential data of relative signal intensity vs electron energy taken when the signals were large.

The boron and titanium pressures were used to obtain the equilibrium constant K for the reaction  $\text{TiB}_2(s) \rightarrow \text{Ti}(g) + 2\text{B}(g)$ . The free energy change was computed ( $\Delta F_T^\circ = -RT \ln K$ ), and the heat of vaporization was obtained from

$$\Delta H_{298}^\circ = \Delta F_T^\circ + T \left( - \frac{\Delta F_T^\circ - \Delta H_{298}^\circ}{T} \right)_{\text{Ti}(s) \rightarrow \text{Ti}(g)} + 2T \left( - \frac{\Delta F_T^\circ - \Delta H_{298}^\circ}{T} \right)_{\text{B}(s) \rightarrow \text{B}(g)} \quad (2)$$

where tabulated<sup>2</sup> free energy functions for Ti and B were used, and free energy functions for  $\text{TiB}_2(s)$  were assumed equal to those for  $\text{Ti}(s) + 2\text{B}(s)$ . This assumption has been shown to be valid up to 700°C by Walker, et al<sup>12</sup>. The heat of formation of  $\text{TiB}_2(s)$  was computed from  $\Delta H_{298}^\circ$  and the tabulated<sup>2</sup> heats of sublimation of boron and titanium.

Data used to compute the heat of vaporization of  $\text{TiB}_2$  are given in Section I of Table I. Ten determinations of  $P_{\text{Ti}}$  relative to  $P_{\text{B}}$  were made over a

temperature range of approximately 200°K. The table gives the equilibrium constant, free energy change, and heat of vaporization of  $\text{TiB}_2(s)$ . The average heat of vaporization at 298°K is  $431.4 \pm 5.9$  kcal/mole, and the heat of formation is  $-53.7 \pm 5.9$  kcal/mole.

TiB<sub>2</sub> in graphite — A crucible fabricated from "National" TSX graphite was used with a  $\text{TiB}_2$  powdered sample. The knife-edged circular orifice in the crucible lid was 0.75 mm in diameter to give  $\sigma_A \leq 5 \times 10^{-3}$ . In a preliminary run the crucible was heated empty to the highest temperatures anticipated for the  $\text{TiB}_2$  measurements to demonstrate that no boron background impurity signals were present. The  $\text{TiB}_2$  sample was then loaded with a 5 mg silver calibration charge and heated to approximately 1100°C where the silver charge was allowed to vaporize completely. This method of calibration has been described by Chupka and Inghram<sup>9</sup>. The pressures of titanium and boron were determined from ion intensity data taken at higher temperatures according to the relation

$$P_x = \frac{I_x T}{K_{\text{Ag}}} \frac{(\sigma i S)_{\text{Ag}}}{(\sigma i S)_x} \quad (3)$$

where  $I_x$  denotes the relative ion current for species x (titanium or boron), T the cell temperature and  $(\sigma i S)$  as described in Equation 1. The constant  $K_{\text{Ag}}$  represents the effective sensitivity constant (the collected charge per effused neutral particle) for the mass spectrometer determined from the silver calibration measurements.

The relative intensities of the boron and titanium signals were observed to increase slowly with time at fixed crucible temperature. Since boron already had the higher partial pressure, this further increase would be paradoxical if titanium and boron only were present. However, equilibration with carbon, while having little effect on the integral free energy for the reaction  $\text{TiB}_2(\text{s}) \rightarrow \text{Ti}(\text{g}) + 2\text{B}(\text{g})$ , can change the partial molal free energies of the titanium and boron markedly. In other terms, the equilibrium constant should remain nearly unchanged when carbon is added while the partial pressures may change significantly.

Equilibrium constants and integral free energies of evaporation were calculated by the Third Law method outlined in equation 2. Ten determinations of  $P_{\text{Ti}}$  and  $P_{\text{B}}$  were made over  $\text{TiB}_2$  in graphite and are given in Table I Section 2. These results yield  $\Delta H_{298}^{\circ} = 429.6 \pm 5.4$  kcal/mole for the vaporization of  $\text{TiB}_2$  and  $\Delta H_{298}^{\circ} = -51.9 \pm 5.4$  kcal/mole for the heat of formation.

Ti+TiB<sub>2</sub> — Excess titanium was added to the graphite crucible containing the  $\text{TiB}_2$  sample used in the runs described above. The crucible was heated for several hours above the melting point of titanium to allow complete reaction, and subsequent X-ray analysis of the sample showed  $\text{TiB}_2$  only, while the inside of the crucible lid showed C and TiC. Additional titanium was added and the heating process was repeated; again the reaction occurred but at much slower rate. A third loading of titanium was added with a weighed amount of silver. A silver calibration was performed at 1100°C and was in agreement (~ 10%) with other calibrations. The vapor pressure of solid titanium was determined from Equation 3: at 1842°K,  $P_{\text{Ti}} = 1.0 \times 10^{-6}$  atm; at 1858°K,  $P_{\text{Ti}} = 1.6 \times 10^{-6}$  atm; and at 1722°K,  $P_{\text{Ti}} = 1.1 \times 10^{-7}$  atm. These results are in excellent agreement with tabulated data<sup>2</sup> and are used to confirm the ionization cross sections computed by Otvos and

Stevenson<sup>11</sup> for titanium. While Ti(s) cannot be in equilibrium with C, TiC and TiB<sub>2</sub>, it is assumed the reaction rate was sufficiently slow after the crucible had been previously exposed to titanium at much higher temperatures to sustain the pressure over solid titanium though the pressure of pure titanium was not sustained over the more reactive liquid. This assumption is corroborated by the observation that even after the second loading of titanium, a plot of  $\ln P_{\text{Ti}}$  vs  $\frac{1}{T}$  taken before the titanium was melted yielded a heat of sublimation of titanium only seven kilocalories larger than given in the JANAF tables.

Below the melting point of titanium the boron signal could not be observed, but at higher temperatures both signals were measured. At the higher temperatures the reaction of titanium again became evident since the vapor pressure (which was determined after each of the three successive titanium loadings) of liquid titanium could not be maintained, although the pressure exceeded that to be expected over TiC + C.

The pressure data are given in Section 3 of Table I and unlike the previous results were not obtained at a fixed position in the phase diagram. As excess titanium was added different phases formed and X-ray analyses showed TiC + C + TiB<sub>2</sub> after the first heating and TiC + TiB (orthorhombic) + TiB<sub>2</sub> after the final heating. The effect of the additional titanium manifests itself in a 20-fold increase in titanium pressure at 2300°K as inspection of the values of  $P_{\text{Ti}}$  in Section 3 of Table I will show and corresponds to the formation of TiB. Titanium diboride was always present and each set of pressures gives consistent values for its heat of vaporization. Results for eleven sets of data give an average heat of vaporization for TiB<sub>2</sub> of 428.6±5.3 kcal/mole and a heat of formation of -50.9±5.3 kcal/mole. The pressures from run No. 10 of Table I taken in the TiC-TiB-TiB<sub>2</sub> region of the phase diagram yield

for the reaction  $\text{TiB(s)} \rightarrow \text{Ti(g)} + \text{B(g)}$ ,  $\log K_{2340} = -11.07$ ,  $\Delta F^\circ_{2340} = 118.5$  kcal/mole, and  $\Delta H^\circ_{298} = 279$  kcal/mole. The associated heat of formation of  $\text{TiB(s)}$  at 298°K is -34 kcal/mole.

Summary — Data from all the experimental points given in Table I are shown in Figure 2. The equilibrium constant for vaporization of  $\text{TiB}_2$  is given as a function of reciprocal temperature. The least-square line through the data gives a heat of vaporization  $\Delta H^\circ_{2287} = 424 \pm 14$  kcal/mole and  $\Delta H^\circ_{298} = 435$  kcal/mole in excellent internal agreement with the Third Law computations.

Searches for Ti-B gaseous species were negative; for example, at 2426°K,  $P_{\text{TiB(g)}}$  was less than  $10^{-9}$  atm during the  $\text{B} + \text{TiB}_2$  runs. Similar bounds for other simple combinations occurred and, in general, extended sweeps through mass 600 indicated no species to ~ 1% of the  $\text{B(g)}$  signal. A search for  $\text{B}_2(\text{g})$  was negative and indicated  $P_{\text{B}_2}$  was less than  $4 \times 10^{-9}$  atm at 2316°K over  $\text{B} + \text{TiB}_2$ . This bound is not low enough to test the tabulated data.<sup>2</sup>

Some impurity signals were observed. In addition to the oxide species which volatilized, elemental  $\text{Fe(g)}$  and  $\text{Cr(g)}$  impurities were observed and never completely disappeared. During the final runs where excess titanium was added the pressure of  $\text{Cr(g)}$  was comparable to  $\text{Ti(g)}$  even though spectroscopic analysis showed the  $\text{Ti(s)}$  to contain no Cr. It is presumed that chromium borides in the  $\text{TiB}_2$  at trace levels were converted to titanium borides and excess chromium. The rather high impurity level of Cr on the inside of the lid corroborates this assumption.

Some unexplained signals occurred at the mass 30 through 36 positions which appeared by every test to be legitimate neutral species leaving the cell during the work with  $\text{B} + \text{TiB}_2$  in  $\text{B}_4\text{C}$ . It is believed they are not related to the Ti-B system but are possibly gaseous B-C species.

In all the previous results the net experimental error assignment comprised the random uncertainties obtained as the standard deviation of the mean compounded with the instrumental uncertainties obtained from the estimated errors in the temperature and vapor pressures. The temperature error has been discussed above; the pressure error was determined from uncertainties in the relative ionization cross sections and in the measured relative multiplier gains for the several species. The internal agreement between the Second and Third Law measurements by Chupka<sup>2</sup> and of the Third Law measurements by Schissel and Williams<sup>3</sup> and of the Second Law data herein presented on the sublimation of boron shows that the relative cross sections  $\sigma_B/\sigma_{Ag}$  predicted by Otvos and Stevenson<sup>11</sup> are in error by less than  $\pm 50\%$ . Assuming the tabulated data<sup>2</sup> on titanium are correct, the pressure determinations on titanium in this study show the relative cross sections  $\sigma_{Ti}/\sigma_{Ag}$  to be in error by less than  $\pm 50\%$ , and thus the ratio  $\sigma_B/\sigma_{Ti}$  is known to at least a factor of two. These observations are consistent with those of Searcy, et al<sup>6</sup> for the free surface vaporization of  $TiB_2$ . The multiplier response for Ag, B, and Ti was measured in this work with an uncertainty of  $\pm 10\%$ . All thermodynamic results taken from tabulated data were assumed to be without error.

## V. DISCUSSION

Partial pressures of titanium and boron have been determined over condensed phases of the Ti-B-C system for several regions of the phase diagram. The first set of measurements was made in the region  $TiB_2$ - $B_4C$ -B, where boron and  $TiB_2$  powders were used in a  $B_4C$  crucible. The second set of measurements was made with  $TiB_2$  powder loaded in a graphite crucible, ostensibly along the line  $TiB_2$ -C in the phase diagram. The third set of measurements was made with titanium metal added to the  $TiB_2$  powder in the graphite crucible used for the



second set of measurements. Results for the heat of vaporization and formation of  $\text{TiB}_2$  are in excellent agreement, giving an average  $\Delta H_{298}^{\circ} = 429.8$  kcal/mole and an average  $\Delta H_{298}^{\circ} = -52.1$  kcal/mole, respectively.

As discussed by Brewer and Haraldsen<sup>1</sup> the observation that  $\text{TiB}_2(\text{s})$  is stable in graphite bounds the stability of  $\text{TiB}_2$  relative to  $\text{TiC}$ . Using the data of Humphrey<sup>13</sup> for  $\text{TiC}$  one finds  $\Delta H_f^{\circ}(\text{TiB}_2) \leq -44$  kcal/mole while the data of Fujishiro and Gokcen<sup>14</sup> would yield  $\Delta H_f^{\circ}(\text{TiB}_2) \leq -31.3$  kcal/mole. Brewer and Haraldsen state that the heats of formation of titanium borides should be about -36 kcal/mole of boron. The agreement between this result and that of Samsonov<sup>5</sup> for  $\text{TiB}_2$  is apparently fortuitous since Samsonov used a value -66 kcal/mole for the heat of formation of  $\text{B}_4\text{C}$  in disagreement with the value used by Brewer and Haraldsen and with recently tabulated values.<sup>2</sup> The present results for  $\text{TiB}_2$  do not agree with  $\Delta H_f^{\circ} = -72$  kcal/mole.

The measurements of Schissel and Williams<sup>3</sup> yield  $\Delta H_f^{\circ} \approx -32$  kcal/mole for  $\text{TiB}_2$ . These mass spectrometric measurements were made with a mixture of boron and  $\text{TiB}_2$  powders contained in a  $\text{TiB}_2$  crucible which was held in a tungsten outer jacket. The boron pressure determinations are in agreement with the data of this report although the titanium pressure was anomalously high. The discrepancy in the titanium pressure is believed to be due to titanium vapor escaping from the region between the  $\text{TiB}_2$  cell and the tungsten jacket, possibly enhanced by reaction with the tungsten, thus leading to erroneous heats of vaporization and formation of  $\text{TiB}_2$ . In a recent experiment with a high temperature calorimeter Lowell and Williams<sup>7</sup> obtained  $\Delta H_f^{\circ} = -50 \pm 5$  kcal/mole for  $\text{TiB}_2$  in agreement with the result of the present investigation.

The question of vaporization coefficients has not been systematically investigated but the values of  $\sigma$  for the two cells were different by one decade and resulted in a statistically insignificant difference in the  $\text{TiB}_2$  values.

The observation that TiB is found with TiB<sub>2</sub> after samples of the appropriate composition range are heated does not preclude the possibility that TiB is stable over a limited temperature range only. However, for TiB(s) to be unstable at low temperatures relative to Ti(s) and TiB<sub>2</sub>(s), the heat of the reaction  $\text{Ti(s)} + \text{TiB}_2\text{(s)} \rightarrow 2\text{TiB(s)}$  must be positive.<sup>15</sup> The value found in this research is -16 kcal/mole. TiB may be unstable at some temperatures relative to decomposition to TiB<sub>2</sub> plus a titanium rich phase, however. If, as suggested by Hansen and Anderkö,<sup>16</sup> TiB decomposes at temperatures above about 2330°K by the reaction  $3\text{TiB(s)} \rightarrow \text{Ti}_2\text{B(s)} + \text{TiB}_2\text{(s)}$ , a few of these measurements may have been made for Ti<sub>2</sub>B and TiB<sub>2</sub> mixtures. However,  $\Delta F$  must be small for the decomposition reaction near the transition point, and no significant error is introduced by the assumption that TiB was the phase present with TiB<sub>2</sub>.

#### ACKNOWLEDGMENT

The authors wish to thank Dr. R. Goton for preparing the B<sub>4</sub>C crucibles and C. E. Lowell for the X-ray analyses. Several helpful discussions with Prof. A. W. Searcy are greatly appreciated.

REFERENCES

1. L. Brewer and H. Haraldsen, J. Electrochem. Soc., 102, 399 (1955).
2. JANAF Interim Thermochemical Tables, The Dow Chemical Co., Midland, Michigan, December 31, 1960.
3. P. O. Schissel and W. S. Williams, Bull. Am. Phys. Soc., Ser II, 4, No. 3 (1959).
4. P. A. Akishin, O. T. Nikitin, and L. N. Gorokov, Doklady Akad. Nauk, S.S.S.R., 129, 1075 (1959).
5. G. V. Samsonov, J. Applied Chem., (USSR) 28, 975 (1955).
6. A. W. Searcy, W. S. Williams, and P. O. Schissel, J. Chem. Phys., 32, 957 (1960).
7. C. E. Lowell and W. S. Williams, Rev. Sci. Instr., in press.
8. M. G. Inghram and R. J. Hayden, National Academy of Sciences and National Research Council, Publication 311 (1954).
9. W. A. Chupka and M. G. Inghram, J. Phys. Chem., 59, 100 (1955).
10. O. C. Trulson, D. E. Hudson, and F. H. Spedding, J. Chem. Phys., in press.
11. J. W. Otvos and D. P. Stevenson, J. Am. Chem. Soc., 78, 546 (1956).
12. B. E. Walker, C. T. Ewing, and R. R. Miller, J. Phys. Chem., 61, 1682 (1957):
13. G. L. Humphrey, J. Am. Chem. Soc., 73, 2261 (1951).
14. S. Fujishiro and N. A. Gokcen, J. Phys. Chem., 65, 161 (1961).
15. A. W. Searcy in Proceedings of Second International High Temperature Symposium, Asilomar, California (Oct. 1959), (McGraw-Hill Book Co., Inc., New York, 1960).
16. M. Hansen and K. Anderko, Constitution of Binary Alloys (McGraw-Hill Book Co., Inc., New York, 1958).

TABLE I

CALCULATION OF THE HEAT OF VAPORIZATION AND  
HEAT OF FORMATION OF  $TiB_2(s)$

System Before Heating.	Run	T (°K)	-log $P_B$	-log $P_{Ti}$	-log K	$\Delta F^\circ$ (kcal/mole)	$\Delta H^\circ_{199}$ (kcal/mole)		$-\Delta H^\circ_f$ (kcal/mole)	
B+ $TiB_2$ in $B_4C$	1	2244		8.47	18.92	194.3	427.5		49.8	
		2296		8.46	18.33	192.6	430.9		53.2	
		2347		8.26	17.62	189.2	432.6		54.9	
		2357		8.44	17.69	190.8	435.2		57.5	
		2382		7.97	16.98	185.0	432.0		54.2	
		2398		7.96	16.81	184.4	432.9		55.2	
		2408		7.74	16.50	181.8	431.2		53.5	
	2	2306		8.40	18.16	191.7	431.1		53.3	
		2462		7.38	15.64	176.2	431.0		53.3	
	3	2411		7.56	16.29	179.7	429.5		61.8	
							431.4			Avg. 53.7±0.7
$TiB_2$ in C	4	2192	6.54	7.19	20.27	203.3	431.4		53.7	
		2185	6.62	7.11	20.36	203.5	430.9		53.2	
	5	2353	5.14	6.31	16.60	178.7	422.7		45.0	
	6	2246	6.20	7.12	19.52	200.6	434.1		56.3	
		2246	6.22	7.00	19.44	199.8	433.2		55.5	
		2246	6.13	7.04	19.30	198.4	431.8		54.1	
		2278	5.78	6.83	18.40	191.8	428.3		50.6	
		2278	5.73	6.78	18.23	190.1	426.6		48.9	
		2278	5.77	6.79	18.33	191.1	427.6		49.9	
		2310	5.57	6.83	17.97	189.9	429.7		52.0	
							429.6			Avg. 51.9±1.1
Ti+ $TiB_2$ in C	7	2271	6.22	6.02	18.46	191.8	427.7		50.0	
		2269	6.18	6.21	18.56	192.7	428.4		50.7	
		2269	6.11	6.29	18.51	192.1	427.9		50.1	
		2307	5.93	6.21	18.06	190.6	430.1		52.4	
		2063	7.61	7.92	23.15	218.5	433.7		56.0	
		2107	7.04	7.59	21.68	209.0	428.6		50.9	
		2272	6.01	6.44	18.45	191.8	427.8		50.1	
	8	2276	5.96	6.55	18.47	192.4	428.8		51.0	
	9	2266	6.61	5.35	18.56	192.5	427.9		50.2	
	10	2340	6.11	4.96	17.18	183.9	426.7		49.0	
		2327	6.05	5.37	17.47	186.0	427.5		49.7	
							428.6			Avg. 50.9±0.7

FIGURE 1

THE CLAUSIUS-CLAPEYRON PLOT FOR  $B^+(g)$ . LOGARITHM OF THE INTENSITY OF  $B^+(g)$  TIMES THE ABSOLUTE TEMPERATURE VS THE RECIPROCAL ABSOLUTE TEMPERATURE.





FIGURE 2

THE TEMPERATURE DEPENDENCE OF THE EQUILIBRIUM  
CONSTANT K OF THE REACTION  $\text{TiB}_2(s) \rightarrow \text{Ti}(g) + 2\text{B}(g)$ .



DISTRIBUTION

Director Advanced Research Projects Agency The Pentagon, Room 3D159 Washington 25, D. C. ATTN: Advanced Propellant Chemistry Office (6)	Commander U. S. Army Ordnance Missile Command Army Rocket and Guided Missile Agency Redstone Arsenal, Alabama ATTN: ORDXR-RFEP (12)
National Aeronautics and Space Administration Washington 25, D. C. ATTN: Office of Technical Information and Educational Programs Code ETL (7)	Commander, U. S. Army Ordnance Missile Command Army Rocket and Guided Missile Agency Redstone Arsenal, Alabama ATTN: ORDXR-RRH Dr. T. A. Barr (1)
National Bureau of Standards Washington 25, D. C. ATTN: Dr. S. Madorskey (1)	Chief, New York Ordnance District U. S. Army 770 Broadway New York 3, New York ATTN: Facilities and Resources Branch (1)
Dr. Charles W. Beckett Chief, Thermodynamic Section Division of Heat and Power National Bureau of Standards Washington 25, D. C. (1)	Chief, New York Ordnance District U. S. Army 770 Broadway New York 3, New York ATTN: ORDEH-O-TA (1)
Commander Oregon Metallurgical Corporation P. O. Box 484 Albany, Oregon ATTN: Mr. Russell G. Hardy (1)	Department of the Army Office, Chief of Ordnance Washington 25, D. C. ATTN: ORDTB (1)
U. S. Atomic Energy Commission Office of Technical Information Extension P. O. Box 62 Oak Ridge, Tennessee (1)	Commanding Officer Ordnance Office Duke Station Durham, North Carolina (1)
U. S. Department of Interior Bureau of Mines 4800 Forbes Street Pittsburgh 13, Pennsylvania ATTN: M. P. Benoy, Reports Librarian Explosives Research Laboratory (1)	Ordnance Materials Research Office Watertown Arsenal Watertown 72, Massachusetts (1)
Commander Army Ballistic Missile Agency Redstone Arsenal, Alabama ATTN: ORDAB-HSI (1)	Dr. A. P. Levitt Chief, High Temperature Branch Watertown Arsenal Laboratories Watertown, Massachusetts (1)
Commander Army Rocket and Guided Missile Agency Redstone Arsenal, Alabama ATTN: Technical Library ORDXR-OTL (4)	Commanding General Aberdeen Proving Ground Maryland ATTN: Ballistic Research Laboratory ORDBG-BLI (1)
Commander U. S. Army Ordnance Missile Command Army Rocket and Guided Missile Agency Redstone Arsenal, Alabama ATTN: ORDXR-OR (1)	Commanding Officer Picatinny Arsenal Dover, New Jersey ATTN: Library (1)
Commander U. S. Army Ordnance Missile Command Army Rocket and Guided Missile Agency Redstone Arsenal, Alabama ATTN: ORDXR-IXD (1)	Commanding Officer Diamond Ordnance Fuze Laboratories Washington 25, D. C. ATTN: ORDTL (012) (1)

Commander U. S. Naval Air Missile Test Center Point Mugu, California ATTN: Technical Library	(1)	Director Special Projects Office Department of the Navy Washington 25, D. C.	(1)
Commander U. S. Naval Ordnance Laboratory White Oak Silver Spring, Maryland ATTN: Library	(1)	Department of the Navy Bureau of Naval Weapons Washington 25, D. C. ATTN: RRRE-6	(1)
Commander U. S. Naval Ordnance Test Station China Lake, California ATTN: Technical Library Branch	(1)	Commander Air Research and Development Center Andrews Air Force Base Washington 25, D. C. ATTN: RDRAPR	(1)
Commanding Officer U. S. Naval Propellant Plant Indian Head, Maryland ATTN: Technical Library	(1)	Commander Air Force Flight Test Center Edwards Air Force Base California ATTN: FTR	(1)
Commander U. S. Naval Weapons Laboratory Dahlgren, Virginia ATTN: Technical Library	(1)	Commander Air Force Flight Test Center Edwards Air Force Base California ATTN: FTRP	(1)
Department of the Navy Office of Naval Research Washington 25, D. C. ATTN: Code 429	(1)	Headquarters 6593rd Test Group (Dev.) Air Force Systems Command Edwards Air Force Base, California ATTN: DGS	(1)
Director U. S. Naval Research Laboratory Washington 20, D. C. ATTN: Chemistry Div., Code 6130 R. R. Miller	(1)	Air Proving Ground Center Elgin Air Force Base Florida ATTN: PGAPI	(1)
Department of the Navy Bureau of Naval Weapons Washington 25, D. C. ATTN: RMMP-2	(1)	Headquarters, Wright Air Development Div. Air Research and Development Command Wright-Patterson Air Force Base Ohio ATTN: WWRCPT-1	(1)
Department of the Navy Bureau of Naval Weapons Washington 25, D. C. ATTN: RMMP-4	(1)	Headquarters Space Systems Division Air Force Systems Command P. O. Box 262, Air Force Unit Post Office Los Angeles 45, California ATTN: TDC	(1)
Department of the Navy Bureau of Naval Weapons Washington 25, D. C. ATTN: RMMP-331	(1)	Commander Aeronautical Systems Division Wright-Patterson Air Force Base Ohio ATTN: ASRCNP	(1)
Department of the Navy Bureau of Naval Weapons Washington 25, D. C. ATTN: DLI-3	(2)	Commander Armed Services Tech. Info. Agency Arlington Hall Station Arlington 12, Virginia ATTN: TIPCR	(10)





British Defence Staff British Embassy 3100 Massachusetts Avenue Washington, D. C. ATTN: Scientific Information Center	(4)	Avco 201 Lowell Street Wilmington, Massachusetts ATTN: Dr. E. Scalla	(1)
Defence Research Member Canadian Joint Staff (W) 2450 Massachusetts Avenue, N. W. Washington 8, D. C.	(4)	Avco 201 Lowell Street Wilmington, Massachusetts ATTN: Dr. H. L. Schick	(1)
Aerojet-General Corporation Box 1947 Sacramento, California ATTN: Technical Information Office	(3)	Battelle Memorial Institute Ceramics Division 505 King Avenue Columbus 1, Ohio ATTN: J. F. Lynch	(1)
Aerojet-General Corporation P. O. Box 296 Azusa, California ATTN: Library	(1)	Battelle Memorial Institute Defense Metals Information Center 505 King Avenue Columbus 1, Ohio ATTN: C. S. Dumont	(1)
Aeronutronic, A Division of Ford Motor Co. Ford Road Newport Beach, California ATTN: L. H. Linder, Manager Technical Information Department	(1)	Borg-Warner Corporation Ingersoll Kalamazoo Division 1810 N. Pitcher Street Kalamazoo, Michigan ATTN: J. W. Schiffel, Chief Engineer Special Projects Department	(1)
Aerospace Corporation P. O. Box 95085 Los Angeles 45, California ATTN: Technical Information Center Acquisitions	(2)	The Carborundum Company Box Number 162 Niagara Falls, New York	(1)
Allied Chemical Corporation General Chemical Division Research Laboratory P. O. Box 405 Morristown, New Jersey ATTN: H. R. Neumart	(1)	Clevite Corporation Mechanical Research Division 540 East 105th Street Cleveland 8, Ohio ATTN: Mr. Gail Davies	(1)
American Cyanamid Company 1937 W. Main Street Stamford, Connecticut ATTN: Dr. A. L. Peiker	(1)	Cornell Aeronautical Laboratories 4455 Genessee Street Buffalo 21, New York ATTN: Mr. J. Beale	(1)
Armour Research Foundation of Illinois Institute of Technology Technology Center Chicago 16, Illinois ATTN: Fluid Dyn. and Prop. Research, Dept. D.	(1)	Cornell University Materials Science Center Thurston Hall Ithaca, New York ATTN: Professor R. L. Sproull	(1)
Atlantic Research Corporation Shirley Highway and Edsall Road Alexandria, Virginia	(1)	DeBell and Richardson, Inc. Hazardville, Connecticut ATTN: Mr. William Eakins	(1)
Atomics International Canoga Park, California ATTN: Mr. J. Hove	(1)	The Dow Chemical Company Security Section Box 31 Midland, Michigan ATTN: Dr. R. S. Karpiuk, 1710 Bldg.	(1)

E. I. duPont deNemours and Company Eastern Laboratory Gibbstown, New Jersey ATTN: Mrs. Alice R. Steward (1)	The Johns Hopkins University Solid Propellant Information Agency Applied Physics Laboratory Silver Spring, Maryland (3)
Esso Research and Engineering Company Chemicals Research Division P. O. Box 51 Linden, New Jersey ATTN: Dr. J. P. Longwell VIA: Chief, New York Ordnance District (1)	The Johns Hopkins University Applied Physics Laboratory 8621 Georgia Avenue Silver Spring, Maryland ATTN: Dr. A. Westenberg (1)
General Electric Company Cincinnati 15, Ohio ATTN: Technical Information Center (1)	Professor J. L. Margrave Chemistry Department University of Wisconsin Madison 6, Wisconsin (1)
Georgia Institute of Technology Mr. Cecil R. Mason, Ceramics Branch Engineering Experiment Station Atlanta, Georgia (1)	Minnesota Mining and Manufacturing Co. 900 Bush Avenue St. Paul 6, Minnesota ATTN: J. W. Millin VIA: R. W. McElroy Security Administrator (2)
The B. F. Goodrich Company Research Center Brecksville, Ohio ATTN: Charles H. Stockman (1)	Northwestern University Materials Research Center The Technological Institute Evanston, Illinois ATTN: Professor M. E. Fine (1)
Grand Central Rocket Company P. O. Box 111 Redlands, California ATTN: Helen Ashman, Librarian (3)	Olin Mathieson Chemical Corporation Research Library I-K-3 275 Winchester Avenue New Haven, Connecticut ATTN: Miss Laura M. Kajuti Mail Control Room (3)
Hercules Powder Company Allegany Ballistics Laboratory P. O. Box 210 Cumberland, Maryland ATTN: Library (2)	Olin Mathieson Chemical Corporation P. O. Box 508 Marion, Illinois ATTN: Research Library (1)
Hercules Powder Company Rocky Hill Plant P. O. Box A Rocky Hill, New Jersey (1)	Rocketdyne 6633 Canoga Avenue Canoga Park, California ATTN: Library 596-306 (3)
Hercules Powder Company Bacchus Works Magna, Utah ATTN: Librarian (1)	Rocketdyne A Division of North American Aviation, Inc. Solid Propulsion Operations P. O. Box 548 McGregor, Texas (1)
Ionics, Inc. 1426 G. Street N. W. Washington 5, D. C. ATTN: Mr. S. G. McGriff Asst. Director of Research (1)	Rocket Power 3016 East Foothill Boulevard Pasadena, California ATTN: Dr. Milton Farber, Vice Pres. (1)
Jet Propulsion Laboratory 4800 Oak Grove Drive Pasadena 3, California ATTN: I. E. Newlan, Chief Reports Group (1)	Shell Development Company 4560 Horton Street Emeryville 8, California (1)

Space Technology Laboratory, Inc. 5730 Arbor-Vitae Street Los Angeles 45, California ATTN: Mr. Robert C. Anderson	(1)	Dr. R. M. Bushong Assistant Technical Director National Carbon Company P. O. Box 560 Niagara Falls, New York	(1)
Thiokol Chemical Corporation Redstone Division Huntsville, Alabama ATTN: Technical Director	(1)	Dr. R. L. Cummerow Parma Research Laboratory P. O. Box 6116 Cleveland 1, Ohio	(1)
Thiokol Chemical Corporation Elkton Division Elkton, Maryland ATTN: Librarian	(1)	Mr. H. A. Downey National Carbon Company Fostoria, Ohio	(1)
Thiokol Chemical Corporation Utah Division Brigham City, Utah ATTN: Library Section	(2)	Dr. D. M. Gillies Union Carbide Metals Co. P. O. Box 580 Niagara Falls, New York	(1)
Thiokol Chemical Corporation Rockets Operation Center P. O. Box 1640 Ogden, Utah ATTN: Librarian	(1)	Mr. D. E. Hamby Union Carbide Corporation 270 Park Avenue Floor 41 New York 17, New York	(1)
United Technology Department P. O. Box 358 Sunnyvale, California ATTN: Librarian	(1)	Dr. A. B. Kinzel Union Carbide Corporation 270 Park Avenue Floor 39 New York 17, New York	(1)
University of Pennsylvania Laboratory for Research on Structure of Matter The Towne Building Philadelphia 4, Pennsylvania ATTN: Professor J. N. Hobstetter	(1)	Haynes Stellite Company Kokomo, Indiana ATTN: Helen F. Kuhns, Librarian	(1)
Wright Aeronautical Corporation Curtiss-Wright Corporation Wood-Ridge, New Jersey	(2)	Dr. R. A. Charpie, Manager Advanced Developments, Union Carbide Corporation 43rd Floor 270 Park Avenue New York 17, New York	(1)
Mr. J. C. Bowman, Director of Research National Carbon Company P. O. Box 6116 Cleveland 1, Ohio	(1)	Dr. R. W. McNamee Union Carbide Corporation 270 Park Avenue 43rd Floor New York 17, New York	(1)
Mr. E. J. Boyle Union Carbide Metals Company P. O. Box 580 Niagara Falls, New York	(1)	Dr. R. M. Milton Linde Company P. O. Box 44 Tonawanda, New York	(1)
Dr. R. G. Breckenridge, Director of Research Parma Research Laboratory P. O. Box 6116 Cleveland 1, Ohio	(1)	Dr. John Pike Parma Research Laboratory P. O. Box 6116 Cleveland 1, Ohio	(1)

Dr. P. O. Schissel  
Parma Research Laboratory  
P. O. Box 6116  
Cleveland 1, Ohio (1)

Dr. Milton Stern  
Speedway Laboratories  
Linde Company  
1500 Polco Street  
Indianapolis 24, Indiana (1)

Mr. W. A. Steiner, Vice Pres. Technology  
National Carbon Company  
270 Park Avenue  
11th Floor  
New York 17, New York (1)

Mr. Roy C. Sundeen  
Parma Research Laboratory  
P. O. Box 6116  
Cleveland 1, Ohio (1)

Mr. J. D. Venables  
Parma Research Laboratory  
P. O. Box 6116  
Cleveland 1, Ohio (1)

Dr. G. H. Wagner  
Union Carbide Corporation  
270 Park Avenue  
7th Floor  
New York 17, New York (1)

Mr. H. L. Willard  
Union Carbide Corporation  
270 Park Avenue  
43rd Floor  
New York 17, New York (1)

UCRI - Library (6)  
Library (1)